

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

2014

A New Species of *Nephus* Mulsant (Coleoptera: Coccinellidae) in Florida

Vince Golia

Wellington, FL, vincegolia@comcast.net

Austin Golia

Wellington, FL

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>

Golia, Vince and Golia, Austin, "A New Species of *Nephus* Mulsant (Coleoptera: Coccinellidae) in Florida" (2014). *Insecta Mundi*. 887. <http://digitalcommons.unl.edu/insectamundi/887>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0372

A New Species of *Nephus* Mulsant (Coleoptera: Coccinellidae) in Florida

Vince Golia and Austin Golia
9841 Scribner Lane
Wellington, FL 33414 USA

Date of Issue: August 29, 2014

Vince Golia and Austin Golia
A New Species of *Nephus* Mulsant (Coleoptera: Coccinellidae) in Florida
Insecta Mundi 0372: 1–3

ZooBank Registered: urn:lsid:zoobank.org:pub:6823DD59-E0F6-4B7E-A939-A1C4A13C08CC

Published in 2014 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Head Layout Editor: Eugenio H. Nearn

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and Submission Requirements available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and AgriFood Canada, Ottawa, ON, Canada
The Natural History Museum, London, UK
Muzeum i Instytut Zoologii PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

A New Species of *Nephus* Mulsant (Coleoptera: Coccinellidae) in Florida

Vince Golia and Austin Golia
9841 Scribner Lane
Wellington, FL 33414 USA
vincegolia@comcast.net

Abstract. A new species of *Nephus* Mulsant (Coleoptera: Coccinellidae) from southern Florida is described: *Nephus* (*Nephus*) *alyssae*. This represents the first member of the subgenus *Nephus* reported in the southeastern United States. Florida species previously placed in the genus *Nephus*, now placed in *Scymnobi* Casey, are compared with the new species.

Introduction

Gordon (1976, 1985) revised *Nephus* Mulsant (Coleoptera: Coccinellidae) for the United States, listing 5 subgenera: *Depressoscymnus* Gordon (1 species), *Nephus* Mulsant (1 species), *Scymnobi* Casey (10 species), *Sidis* Mulsant (1 species), and *Turboscymnus* Gordon (1 species). Peck and Thomas (1998) followed this arrangement in their checklist of Florida beetles, listing only three species of *N.* (*Scymnobi*) as occurring in Florida. Gordon and González (2002) raised *Scymnobi* to full generic rank. Specimens of an unknown species were recently collected in Florida and one was sent to Robert D. Gordon for identification. He identified it as an undescribed species of *Nephus*. The only other member of *Nephus* (*Nephus*) in North America is *Nephus ornatus* (LeConte) occurring in the northern US. The new species is described and compared with *N. ornatus* and similar appearing members of *Scymnobi* in Florida.

Materials

Specimens studied are deposited in the following collections: **ABSC**—Archbold Biological Station, Venus, Florida; **FSCA**—Florida State Collection of Arthropods, Gainesville, Florida; **USNM**—United States National Museum, Smithsonian Institution, Washington, District of Columbia; **VGIC**—Personal collection, Vince Golia, Wellington, Florida.

Nephus (*Nephus*) *alyssae* Golia and Golia, n. sp.

Figures 1, 4, 5

Diagnosis. The small size, elytron with single golden central spot, and dorso-ventrally flattened body distinguishes *N. alyssae* from any other US species of *Nephus*.

Description. Scymnini. Female. Body 1.47–1.87 mm long; width 0.88–1.15 mm (holotype length 1.65 mm; width 1.00 mm). Body oval, elongate (Fig. 1), dorsoventrally flattened (Fig. 5); entirely dark brown to black with head and legs brownish-yellow and each elytron with single golden central spot ranging from 0.66–0.95 mm on apical 2/3; dorsum of body covered with yellow and white, short semi-erect hairs. Head not concealing prosternum; antennae 11-segmented with a small, symmetrical club. Prosternum lacking coxal lines, flattened, punctate. Abdominal sternite I with post coxal lines long, curving forward laterally, but not attaining lateral or basal margins. Female spermatheca shortened, truncate (Fig. 4), similar to that of *N. ornatus* (Gordon 1985). Male unknown.

Distribution. Known only from south Florida.

Type material. **Holotype**, female with label data “Florida, Palm Beach Co., Lake Worth, Hypoluxo, Hypoluxo Scrub N.A.; November 10, 2009; Vince Golia; ‘sweeping’, 26.566642,-80.056759” deposited in FSCA.

Paratypes. Nine females total: Florida. Palm Beach County. Lake Worth, Hypoluxo Road & Edisto Drive. August 29, 2006, Vince Golia; 'sweeping', 26.571885,-80.161352 (1 female; USNM). FLORIDA. Palm Beach County. Lake Worth, Hypoluxo Road & Lyons Road, September 16, 2011, Vince Golia; 'sweeping', 26.568505,-80.191304. (1 female; FSCA). Florida, Broward Co., Fort Lauderdale, Macintosh Road; July 12, 2013; Stephen Beidler; 'sweeping', 26.08179,-80.126356 (2 females; FSCA). Florida, Broward Co., Fort Lauderdale, Macintosh Road; October 17, 2013, Vince Golia; 'sweeping', (26.077443,-80.124391) (4 females; ABSC; FSCA; VGIC). Florida. Dade County, Miami. NW 2nd Avenue & NW 161 Street. April 10, 2010, Vince Golia; 'sweeping', 25.922244,-80.203580, (1 teneral female; FSCA).

Remarks. In North American *Nephus* and *Scymnobi*, the shortened truncate female spermatheca is similar only to that of *N. ornatus*. Even without males, *N. alyssae* is readily distinguishable from all North American species of Scymnini. *Nephus alyssae*, with a single spot on each elytron, is readily distinguished from *N. ornatus* which has two spots on each elytron. In the US, *Scymnobi* species are not dorsoventrally flattened (Fig. 6), and are easily distinguished from *N. alyssae* based on that character alone. In comparison with other Florida species of *Scymnobi*, *S. bivulnerus* (Horn) has one red spot on each elytron and red a pronotum and *S. intrusus* (Horn) is completely brown in color. *Scymnobi flavifrons* (Melsheimer) can be confused with *N. alyssae*, and can be found at the same locality. In *S. flavifrons*, the body is not dorsoventrally flattened and the golden to reddish-orange elytral spots vary from a small spot near the apex (Fig. 3) to occupying 2/3 of the elytra (Fig. 2) as in *N. alyssae*. However, the spots in *N. alyssae* are always the same size and golden, and the body is distinctly flattened (Fig. 1, 5).

All ten specimens of *Nephus alyssae* were collected while sweeping plants along roadsides where the land is disturbed in three counties known to have many exotic species. As with *Nephus (Sidis) binaevatus* (Mulsant) in California (Gordon 1976, 1985), *N. alyssae* may yet be recognized as an immigrant species. It is believed that all *Nephus* spp. are predators of mealy bugs.

Etymology. We name this species after my daughter and Austin's sister, who fondly remembers the joy of finding her first lady bug. Naming this species for Alyssa will allow her to remember that joy forever.



Figures 1-6. 1) *Nephus alyssae*, dorsal habitus. 2) *Scymnobi flavifrons*, dorsal habitus, Archbold Biological Station, Highlands Co., FL. 3) *Scymnobi flavifrons*, dorsal habitus, Lake Worth, Palm Beach Co., FL. 4) *Nephus alyssae*, spermatheca. 5) *Nephus alyssae*, lateral habitus. 6) *Scymnobi flavifrons*, lateral habitus, Archbold Biological Station.

Acknowledgments

Many thanks to Paul Skelley (FSCA), Mike Thomas (FSCA), Ian Stock (FSCA), Mark Deyrup (ABSC), and Robert Gordon (Northern Plains Entomology, Willow City, ND) for their time and expertise. We thank Andrew Cline, California State Collection of Arthropods, Sacramento, California, for loan of *N. binaevatus*. We thank Mark Deyrup, Archbold Biological Station, and Robert Gordon, Northern Plains Entomology, for reviews of the manuscript.

Literature Cited

- Gordon, R. D. 1976.** The Scymnini (Coleoptera: Coccinellidae) of the United States and Canada: key to genera and revision of *Scymnus*, *Nephus* and *Diomus*. Bulletin of the Buffalo Society of Natural Sciences 28: 1–362.
- Gordon, R. D. 1985.** The Coccinellidae (Coleoptera) of America north of Mexico. Journal of the New York Entomological Society 93: 1–912.
- Gordon, R. D., and G. F. González. 2002.** South American Coccinellidae (Coleoptera). Part IX: a systematic revision of *Scymnobioides* Casey (Scymninae: Scymnini). Frustula Entomologica 25(38): 57–85.
- Peck, S. B., and M. C. Thomas. 1998.** A distributional checklist of the beetles of Florida. Arthropods of Florida and Neighboring Land Areas 16: 1–180.

Received July 17, 2014; Accepted August 20, 2014
Review Editor Paul Skelley

